



Contribution ID: 131

Type: Talk

## Dispersive High-Energy Spin Excitations In Iron Pnictide Superconductors Investigated With Resonant Inelastic X-Ray Scattering

*Saturday, 17 September 2011 14:50 (20 minutes)*

The discovery of iron-based high temperature superconductivity has triggered tremendous research efforts in searching for novel high-Tc superconductors and in understanding the related fundamental physics. Unlike the cuprates whose parent compounds are long-range ordered antiferromagnetic Mott insulators, the iron-based parent compounds are 'spin-density wave' metals with delocalized electronic structure and more itinerant magnetism. Recent ARPES studies provides cumulative evidence that superconductivity in iron-based materials may be connected with interband pair scattering between the quasi-nested electron-hole Fermi surfaces. On the other hand, the observation of spin fluctuations by Inelastic Neutron Scattering (INS) in these materials, similar to those seen in cuprates, suggests that cuprate and iron-based high-Tc superconductors may share a common pairing mechanism.

Recent developments of the high-resolution resonant inelastic X-ray scattering (RIXS) technique [1] have enabled investigations of magnetic excitations in cuprates [2,3], which show excellent agreement with results from INS. In this presentation we demonstrate that RIXS can be used to measure collective magnetic excitations in iron-based superconductors despite their much stronger itinerancy compared to cuprates. The persistence of high-energy spin excitations even in optimally doped pnictide superconductors in a wide range of temperatures strongly suggests a spin-mediated Cooper pairing mechanism as proposed in cuprate superconductors.

[1] G. Ghiringhelli et al., Rev. Sci. Instrum. 77, 113108 (2006); V. N. Strocov et al., J. Synch. Radiat. 17, 631 (2010)

[2] J. Schlappa et al., Phys. Rev. Lett. 103, 047401 (2009)

[3] L. Braicovich et al., Phys. Rev. Lett. 104, 077002 (2010)

### Please specify the session

RIXS

### Please specify poster or talk

Talk

**Primary author:** Dr ZHOU, K J (Swiss Light Source, Paul Scherrer Institut, CH 5232 Villigen PSI, Switzerland)

**Co-authors:** Dr MONNEY, C (Swiss Light Source, Paul Scherrer Institut, CH-5232 Villigen PSI, Switzerland); Dr DING, H (Beijing National Laboratory for Condensed Matter Physics, and Institute of Physics, Chinese Academy of Sciences, Beijing 100190, China); Dr VAN DEN BRINK, J (Institute for Theoretical Solid State Physics, IFW Dresden, 01171 Dresden, Germany); Dr WANG, N L (Beijing National Laboratory for Condensed Matter Physics, and Institute of Physics, Chinese Academy of Sciences, Beijing 100190, China); Dr DAI, P C (Beijing National Laboratory for Condensed Matter Physics, and Institute of Physics, Chinese Academy of Sciences, Beijing 100190, China and Department of Physics and Astronomy, The University of Tennessee, Knoxville, Tennessee 37996, USA)

and Neutron Scattering Sciences Division, Oak Ridge National Laboratory, Oak Ridge, Tennessee 37831, USA); Dr SCHMITT, T (Swiss Light Source, Paul Scherrer Institut, CH-5232 Villigen PSI, Switzerland); Dr STROCOV, V N (Swiss Light Source, Paul Scherrer Institut, CH-5232 Villigen PSI, Switzerland); Dr DAI, X (Beijing National Laboratory for Condensed Matter Physics, and Institute of Physics, Chinese Academy of Sciences, Beijing 100190, China); Dr HUANG, Y B (Beijing National Laboratory for Condensed Matter Physics, and Institute of Physics, Chinese Academy of Sciences, Beijing 100190, China and Swiss Light Source, Paul Scherrer Institut, CH 5232 Villigen PSI, Switzerland)

**Presenter:** Dr ZHOU, K J (Swiss Light Source, Paul Scherrer Institut, CH 5232 Villigen PSI, Switzerland)

**Session Classification:** Resonant Inelastic and Elastic X-ray Scattering

**Track Classification:** Resonant Inelastic and Elastic X-ray Scattering