


Challenges and perspectives in resonator-mediated quantum many-body physics: From atoms to solid state



Contribution ID: 2

Type: not specified

”Engineering interactions in the Quantum Hall effect through vacuum field in metamaterial cavities”

Monday, 17 June 2024 09:00 (1 hour)

Jérôme Faist¹, J. Enkner¹, L. Graziotto¹, M. Beck¹, C. Reichl², W. Wegscheider², G. Scalari¹

¹ Institute of Quantum Electronics, ETH Zurich

² Laboratory for Solid State Physics, ETH Zurich, Zurich 8093, Switzerland

In a microcavity, the strength of the electric field caused by the vacuum fluctuations, to which the strength of the light-matter coupling WR is proportional, scales inversely with the cavity volume. One very interesting feature of the circuit-based metamaterials is the fact that this volume can be scaled down to deep subwavelength values in all three dimension of space.¹ We have used transport to probe the ultra-strong light-matter coupling² and shown that the latter can induce a breakdown of the integer quantum Hall effect³. The phenomenon is explained in terms of cavity-assisted hopping, an anti-resonant process where an electron can scatter from one edge of the sample to the other by “borrowing” a cavity photon from the vacuum⁴. Recently a proposal suggested that the value of the quantized Hall voltage can be renormalized by the cavity⁵, but later work demonstrated that such renormalization corresponds to a singular point in the parameter space. We have investigated this effect experimentally using a Wheatstone bridge geometry⁶ and found the quantization to be held⁷.

We have also investigated a new experimental geometry where a hovering resonator is positioned with nanoactuators above the Hall bar, providing a way to continuously vary the light-matter coupling up to a value of $WR/w=0.33$ while the sample is maintained at millikelvin temperatures. Using this approach, we observe the effect of light-matter coupling on the effective electron g -factor as well as its effect on the gap of the Laughlin states. In particular, we observe an enhancement of the $5/3, 4/3$ and $5/7$ fractional state gaps by up to a factor of two⁸.

1. Scalari, G. et al. Ultrastrong Coupling of the Cyclotron Transition of a 2D Electron Gas to a THz Metamaterial. *Science* 335, 1323–1326 (2012).

2. Paravicini-Bagliani, G. L. et al. Magneto-Transport Controlled by Landau Polariton States. *Nat. Phys.* 15, 186–190 (2019).

3. Appugliese, F. et al. Breakdown of topological protection by cavity vacuum fields in the integer quantum Hall effect. *Science* 375, 1030–1034 (2022).

4. Ciuti, C. Cavity-mediated electron hopping in disordered quantum Hall systems. *Phys. Rev. B* 104, 155307

(2021).

5. Rokaj, V., Penz, M., Sentef, M. A., Ruggenthaler, M. & Rubio, A. Polaritonic Hofstadter butterfly and cavity control of the

quantized Hall conductance. *Phys. Rev. B* 105, 205424 (2022).

6. Schopfer, F. & Poirier, W. Testing universality of the quantum Hall effect by means of the Wheatstone bridge. *J. Appl. Phys.*

102, 054903 (2007).

7. Enkner, J. et al. Testing the Renormalization of the von Klitzing Constant by Cavity Vacuum Fields. Preprint at

<https://doi.org/10.48550/arXiv.2311.10462> (2023).

8. Enkner, J. et al. Enhanced fractional quantum Hall gaps in a two-dimensional electron gas coupled to a hovering split-ring

resonator. Preprint at <https://doi.org/10.48550/arXiv.2405.18362> (2024).

Presenter: FAIST, Jérôme (ETH Zurich)