

Low temperature magnetic structure and lattice anomalies at the commensurate-incommensurate transition of multiferroic

Thursday, 19 September 2013 12:30 (2 hours)

Ferroelectric materials have been known since almost one century ago [1]. While their potential for applications was rapidly recognized, the possibility of combining ferroelectricity with magnetic order -preferably with ferromagnetism- has resulted in an enormous deal of interest during the last decade. Several new materials combining both types of order have been reported, although their promising multifunctionalities have been obscured by two facts: one one side, most of them are antiferromagnetic. On the other, their transition temperatures are too low for most practical applications (typically below 40K).

The oxygen-deficient double perovskite YBaFeCuO5 constitutes a remarkable exception. Spontaneous electric polarization has been recently reported to exist below an unusually high temperature of $T_C \approx 230\text{K}$ [2] coinciding with the occurrence of a commensurate - to - incommensurate reorientation of the Fe³⁺ and Cu²⁺ magnetic moments [3,4]. From a more fundamental point of view the observation of incommensurable magnetic order in a tetragonal material at such high temperatures is rather surprising. In particular, the nature of the relevant competing magnetic interactions and its possible link to low dimensionality or geometrical frustration is not understood at present.

Although the existence of the spin reorientation in this material is know since 1995 [3] the low temperature magnetic structure has not yet been solved. Using neutron powder diffraction we have recently been able to propose a spiral model which satisfactorily describes the measured magnetic intensities below T_C [4]. Also, investigation of the crystal structure showed the existence of small anomalies in the lattice parameters and some interatomic distances at T_C . The relevance of these findings for the magnetoelectric coupling, the direction of the polarization, the modification of the different exchange paths in the structure and the stabilization of the incommensurate magnetic order below T_C are discussed.

- [1] J. Valasek Physical Review 15, 537(1920)
- [2] B. Kundys et al., Appl. Phys. Lett. 94, 072506, (2009).
- [3] V. Caignaert et al., J. Solid State Chem. 114, 24, (1995).
- [4] M. Morin et al., in preparation

Primary author: Mr MORIN, Mickael (Paul Scherrer Institut)

Co-authors: Dr SCARAMUCCI, Andrea (Materials Theory, ETH Zurich, Wolfgang-Pauli-Strasse 27, CH-8093 Zurich, Switzerland); CHEPTIAKOV, Denis (Paul Scherrer Institut); Dr POMJAKUSHINA, Ekaterina (Laboratory for Developments and Methods, Paul Scherrer Institut, 5232 Villigen PSI, Switzerland); Dr DENG, Guochu (Bragg Institute, ANSTO, New Illawarra Road, Lucas Height, New South Wales 2233, Australia); Dr RODRÍGUEZ-CARVAJAL, Juan (Institut Laue-Langevin, BP 156X, F-38042 Grenoble, France); Prof. CONDER, Kazimierz (Laboratory for Developments and Methods, Paul Scherrer Institut, 5232 Villigen PSI, Switzerland); Dr KELLER, Lukas (Laboratory for Neutron Scattering, Paul Scherrer Institut, 5232 Villigen PSI, Switzerland); Dr MEDARDE, Marisa (Paul Scherrer Institut); KENZELMANN, Michel (Paul Scherrer Institut); Prof. SPALDIN, Nicola (Materials Theory, ETH Zurich, Wolfgang-Pauli-Strasse 27, CH-8093 Zurich, Switzerland)

Presenter: Mr MORIN, Mickael (Paul Scherrer Institut)

Session Classification: Poster session II and lunch