

Neutron Polarimetry On the 10^{-4} Level

Tuesday 10 September 2013 16:45 (15 minutes)

Correlation coefficients of the polarized neutron beta decay offer a way to test the standard model by precision measurements. One of the currently leading systematic errors is the neutron polarization. State-of-the-art polarizing super mirrors in the X-SM geometry deliver about 99.7(1)% polarization.

We present recent developments in cold neutron polarization, based on the opaque test bench of ^3He spin filters: We have shown experimentally that the accuracy of polarization analysis with opaque ^3He spin filters is better than 10^{-4} . By optimizing the operating conditions of supermirrors and selecting the materials, we have achieved a polarization of 99.97(1) % with the X-SM geometry for a divergent 5.3 Å beam (monochromatized by a velocity selector). These results solve the issue of neutron beam polarization for the next generation of neutron beta decay correlation experiments which aim for accuracies of 10^{-4} .

Primary author: KLAUSER, Christine (Institut Laue-Langevin, Atominstitut TU Wien)

Co-authors: JULLIEN, David (Institut Laue-Langevin); CHASTAGNIER, Jérémie (Institut Laue-Langevin); SCHNEIDER, Martin (SwissNeutronics); REBROVA, Nataliya (Universität Heidelberg); BÖNI, Peter (TU München); GUIMER-A-MILLÀN, Pilar (Institut Laue-Langevin); BIGAULT, Thierry (Institut Laue-Langevin); SOLDNER, Torsten (Institut Laue-Langevin)

Presenter: KLAUSER, Christine (Institut Laue-Langevin, Atominstitut TU Wien)

Session Classification: Tu - 4