

## Search for LFV and rare decays at the NA62 experiment at CERN

*Tuesday 10 September 2013 18:00 (3 hours)*

The decay  $K^+ \rightarrow \pi^+ \nu \bar{\nu}$  is highly suppressed in the Standard Model (SM), while its rate can be predicted with minimal theoretical uncertainty. The branching ratio for this decay is thus a sensitive probe of the flavor sector of the SM.; however, the smallness of this BR ( $8 \times 10^{-11}$ ) and challenging experimental signature make it very difficult to measure. The primary goal of the NA62 experiment at the CERN SPS is to measure  $BR(K^+ \rightarrow \pi^+ \nu \bar{\nu})$  with  $\sim 10\%$  precision. This will require the observation of  $10^{13}$   $K^+$  decays in the experiment's fiducial volume, as well as the use of high-performance systems for precision tracking, particle identification, and photon vetoing. These aspects of the experiment will also allow NA62 to carry out a rich program of searches for lepton flavor and/or number violating  $K^+$  decays. Such searches can probe new physics scenarios involving, for example, heavy Majorana neutrinos or R-parity violating SUSY. Part of the experimental apparatus was commissioned during a technical run in 2012; installation continues and data taking is expected to begin in late 2014. The physics prospects and the status of the NA62 experiment will be reviewed.

**Primary author:** LAZZERONI, Cristina (University of Birmingham, UK)

**Presenter:** LAZZERONI, Cristina (University of Birmingham, UK)

**Session Classification:** Poster, BBQ & Drinks