

Search for LFV and rare decays at the NA62 experiment at CERN

Tuesday 10 September 2013 18:00 (3 hours)

The decay $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ is highly suppressed in the Standard Model (SM), while its rate can be predicted with minimal theoretical uncertainty. The branching ratio for this decay is thus a sensitive probe of the flavor sector of the SM.; however, the smallness of this BR (8×10^{-11}) and challenging experimental signature make it very difficult to measure. The primary goal of the NA62 experiment at the CERN SPS is to measure $BR(K^+ \rightarrow \pi^+ \nu \bar{\nu})$ with $\sim 10\%$ precision. This will require the observation of 10^{13} K^+ decays in the experiment's fiducial volume, as well as the use of high-performance systems for precision tracking, particle identification, and photon vetoing. These aspects of the experiment will also allow NA62 to carry out a rich program of searches for lepton flavor and/or number violating K^+ decays. Such searches can probe new physics scenarios involving, for example, heavy Majorana neutrinos or R-parity violating SUSY. Part of the experimental apparatus was commissioned during a technical run in 2012; installation continues and data taking is expected to begin in late 2014. The physics prospects and the status of the NA62 experiment will be reviewed.

Primary author: LAZZERONI, Cristina (University of Birmingham, UK)

Presenter: LAZZERONI, Cristina (University of Birmingham, UK)

Session Classification: Poster, BBQ & Drinks