



Contribution ID: 335

Type: Oral presentation

Timing of Holocene volcanic eruptions and their radiative aerosol forcing

Thursday, 6 October 2022 14:50 (20 minutes)

Volcanic eruptions play a dominant role in driving climate, in ways beyond the established short-term influence on surface air temperatures. In order to mitigate and adapt to the climate effects of future large volcanic eruptions we need to better quantify the risk of these eruptions including (a) the probability of their occurrence and (b) their expected climatic impact. The observational record of the timing of volcanic eruptions, their locations, magnitudes of sulphate aerosol injection is incomplete which limits our understanding of the sensitivity of the Earth system to volcanism and the vulnerability of social and economic systems to the climate impact of past and future eruptions.

Here we use an array of synchronized, accurately dated, high-resolution ice-core aerosol records from Greenland and Antarctica to reconstruct the timing, sulphur injections and source locations of 850 volcanic eruptions occurring during the Holocene (i.e., the past 11,500 years) to answer the questions:

- 1) What is the likelihood of a stratospheric sulphur injection as large as that from the colossal eruption of Tambora in 1815 to occur somewhere on the globe within the next 50 years?
- 2) How has subaerial volcanic activity changed in space and time throughout the Holocene?
- 3) How did major eruptions affect global climate and humans through time?

We demonstrate in case studies (e.g., McConnell et al., 2020; Pearson et al., 2022) how novel geochemical tools (e.g., sulphur isotopes, cryptotephra) allow to constrain source parameters of the eruptions (location, plume injection height, stratospheric vs. tropospheric aerosol formation) that control their effects on climate, and propose exact dates for the largest eruptions of the Holocene with the help of ultra-long tree-ring chronologies.

Finally, we generate global-scale, space-and-time resolved stratospheric aerosol properties for climate models (HolVol1.0; Sigl et al., 2021; Abbott et al., 2021) to simulate the volcanic influence on Holocene climate evolution and examine to which extent simulated climate responses agree with those inferred from proxy records.

References:

Abbott, P. M. et al. Volcanic climate forcing preceding the inception of the Younger Dryas: Implications for tracing the Laacher See eruption. *Quaternary Sci Rev* 274, (2021).

McConnell, J. R. et al. Extreme climate after massive eruption of Alaska's Okmok volcano in 43 BCE and effects on the late Roman Republic and Ptolemaic Kingdom. *P Natl Acad Sci USA* 117, 15443-15449, doi:10.1073/pnas.2002722117 (2020).

Pearson, C. et al. Geochemical ice-core constraints on the timing and climatic impact of Aniakchak II (1628 BCE) and Thera (Minoan) volcanic eruptions. *P Natl Acad Sci USA Nexus* (2022).

Sigl, M., et al. Volcanic stratospheric sulfur injections and aerosol optical depth during the Holocene (past 11,500 years) from a bipolar ice core array. *Earth Syst. Sci. Data* 2022, 1-45, doi:10.5194/essd-2021-422 (2022).

Primary authors: SIGL, Michael (OCCR/University of Bern); Dr ABBOTT, Peter (OCCR/University of Bern); Dr ADOLPHI, Florian (Alfred-Wegener-Institut Helmholtz-Zentrum für Polar- und Meeresforschung); Dr ALBERT,

Paul G. (Department of Geography, College of Science, Swansea University); CHELLMAN, Nathan (Desert Research Institute); Prof. BAILLIE, Michael G. L. (School of Natural and Built Environment, Queen's University Belfast, UK.); BURKE, Andrea (University of St Andrews); COLE-DAI, Jihong (South Dakota State University); COOK, Eliza (Centre for Ice and Climate, Niels Bohr Institute, University of Copenhagen); Prof. DAVIES, Siwan (Department of Geography, College of Science, Swansea University); GABRIEL, Imogen (OCCR/University of Bern); HELMICK, Meredith (Climate Change Institute, University of Maine); Dr HÖRHOLD, Maria (Alfred-Wegener-Institut Helmholtz-Zentrum für Polar- und Meeresforschung); Ms INNES, Helen (School of Earth and Environmental Sciences, University of St Andrews); Prof. JUNGCLAUS, Johann (Max-Planck Institut für Meteorologie,); Prof. KRÜGER, Kirstin (Department of Geosciences, University of Oslo); Prof. NICOLUSSI, Kurt (Department of Geography, University of Innsbruck); Prof. KURBATOV, Andrei (Climate Change Institute, University of Maine); Mrs LIN, Jiamei (University of Copenhagen); Prof. MCCONNELL, Joseph (Desert Research Institute); Prof. PEARSON, Charlotte (Laboratory of Tree-Ring Research, University of Arizona); Dr PLUNKETT, Gill (School of Natural and Built Environment, Queen's University Belfast); Dr SALZER, Matthew (Laboratory of Tree-Ring Research, University of Arizona); SEVERI, Mirko (Department of Chemistry Ugo Schiff, University of Florence); SVENSSON, Anders (Physics of Ice, Climate and Earth, Niels Bohr Institute, University of Copenhagen); Prof. TIMMRECK, Claudia (Max-Planck Institut für Meteorologie); Dr TOOHEY, Matthew (Institute of Space and Atmospheric Studies, Department of Physics & Engineering Physics, University of Saskatchewan)

Presenter: SIGL, Michael (OCCR/University of Bern)

Track Classification: Holocene and last 2000 year climate forcings and variability