



Abstract ID : 173

# Understanding the heterogeneous global climate imprint of Dansgaard-Oeschger Events and Heinrich Events - categorizing the events

## Content

Jonathan Ortved Melcher 1, Sune Halkjær 1, S. O. Rasmussen 1, and G. Vettoretti 1

1 Physics of Ice, Climate and Earth, Niels Bohr Institute, University of Copenhagen, Tagensvej 16, 2200 Copenhagen, Denmark

The climate of the last glacial period was dominated by abrupt Dansgaard-Oeschger Events where climate changed between mild and cold phases, probably related to changes in the Atlantic Ocean circulation. During some cold phases, ice sheets formed massive armadas of icebergs which further influenced both the ocean and atmosphere. In some areas, the climate response of Heinrich and Dansgaard-Oeschger Events are more and less extreme versions of the same pattern, but as we move further away from the North Atlantic and downwind areas, the response becomes more heterogeneous.

In a series of three contributions (by Rasmussen, Halkjær, and Melcher, respectively), we explore the heterogeneous global climate imprint of Dansgaard-Oeschger Events and Heinrich Events using a combination of palaeoclimate data and model data from a version of the CCSM4 model which exhibits self-sustained (i.e. unforced) Dansgaard-Oeschger Events, on top of which we forced Heinrich-type freshwater events. The dynamics of Dansgaard-Oeschger Events has previously been analyzed using a 3x3-degree version of the model in Vettoretti et al [2022], where it was shown that the atmospheric CO<sub>2</sub> content controls whether the model operates in an oscillatory regime as well as how CO<sub>2</sub> controls the duration of the mild interstadial periods. Here, we use a 1x1-degree model run similar to that of Vettoretti and Peltier [2018] but apply freshwater forcing to selected stadial periods to emulate the effect of Heinrich Events. We compare the model data to the speleothem data set gathered by Corrick et al. [2020] and selected ocean-sediment and ice-core records.

We aim to develop statistical criteria to categorize which areas in the model show climate changes resembling (1) Dansgaard-Oeschger Events, (2) Heinrich events, (3) Antarctic AIM events, and combinations thereof. The poster discusses the criteria employed and presents examples.

### References

- Corrick et al., Synchronous timing of abrupt climate changes during the last glacial period, *Science* vol. 369 (6506), pp. 963-969, 2020, doi: 10.1126/science.aay5538
- Vettoretti and Peltier, Fast Physics and Slow Physics in the Nonlinear Dansgaard-Oeschger Relaxation Oscillation, *J. Climate* vol. 31, pp. 3423-3449, 2018, doi: 10.1175/JCLI-D-17-0559.1
- Vettoretti et al., Atmospheric CO<sub>2</sub> control of spontaneous millennial-scale ice age climate oscillations, *Nature Geoscience* vol. 15, pp. 300-306, 2022, doi: DOI 10.1038/s41561-022-00920-7

**Primary author:** Mr MELCHER, Jonathan Ortved (Physics of Ice, Climate and Earth, Niels Bohr Institute, University of Copenhagen.)

**Co-authors:** Mr HALKJÆR, Sune (Physics of Ice, Climate and Earth, Niels Bohr Institute, University of Copenhagen.); Prof. OLANDER RASMUSSEN, Sune (Physics of Ice, Climate and Earth, Niels Bohr Institute, University of Copenhagen. ); Dr VETTORETTI, Guido (Physics of Ice, Climate and Earth, Niels Bohr Institute, University of Copenhagen.)

**Presenter:** Mr MELCHER, Jonathan Ortved (Physics of Ice, Climate and Earth, Niels Bohr Institute, University of Copenhagen.)

**Track Classification:** Rapid changes and teleconnections