



Contribution ID: 22

Type: **Invited**

Quantitative simulation tools for predicting radiation damage driven by high-intensity x-ray pulses

Friday, 16 October 2020 13:30 (20 minutes)

One of the key opportunities offered by the development of x-ray free-electron lasers is the determination, at atomic resolution, of the three-dimensional structure of biologically relevant macromolecules. The basic idea underlying molecular imaging using x-ray free-electron lasers is the “diffract-and-destroy” concept: If one uses an x-ray pulse that is sufficiently short (on the order of femtoseconds), then in a single shot an x-ray scattering pattern may be obtained that is practically unaffected by atomic displacements triggered by ionization events during the x-ray pulse. What cannot be eliminated in this way is the impact of the electronic damage on the x-ray scattering patterns. Theory, therefore, plays an important role in the development of this new imaging technique: A quantitative understanding is required of the damage processes occurring during the exposure of a molecule to an ultraintense, ultrafast x-ray pulse. In this talk, I will present progress we have made in order to address this challenge. One tool we have developed, XMDYN, is a molecular-dynamics code that utilizes ab-initio atomic electronic-structure information, computed on the fly, within a Monte-Carlo framework. XMDYN has been successfully tested through experiments at LCLS and SACLA. XMDYN is part of a powerful start-to-end simulation framework for single-particle imaging at the European XFEL. Recently, we have taken first steps towards a full ab-initio framework for simulating high-intensity x-ray-matter interactions. Our new XMOLECULE software solves the polyatomic quantum-mechanical electronic-structure problem for every electronic state arising during the exposure of a molecule to a strong x-ray pulse. From this information, electronic transition rates (such as Auger decay rates) are computed on the fly, and the associated rate equations are integrated utilizing a Monte-Carlo method. XMOLECULE played a key role in a recent LCLS experiment on iodomethane, in which hard x-rays focused to a peak intensity exceeding 10^{19} W/cm² produced the highest charge states ever formed using light. Not only did XMOLECULE correctly predict the charge-state distribution observed, but it also helped identify a new molecular ionization enhancement mechanism based on intramolecular charge transfer.

Primary author: SANTRA, Robin (DESY & Universität Hamburg)

Presenter: SANTRA, Robin (DESY & Universität Hamburg)

Session Classification: Session 4 - Radiation Damage in Complementary Fields including Biological Imaging